



A Review of *Barleria prionitis* L. a Rarely Known Plant with Potential of Phytochemical and Pharmacological Assets

**Vaishnavi V. Khandre ^a, Pallavi L. Bansod ^a,
Fulchand V. Kajale ^{a*}, N. B. Ghiware ^{b++} and A. B. Roge ^c**

^a Department of Pharmacology, Nanded Pharmacy College, Nanded. Swami Ramanand Teerth Marathwada University, Nanded, India.

^b Nanded Pharmacy College, Nanded. Swami Ramanand Teerth Marathwada University, Nanded, India.

^c Department of Pharmaceutical Chemistry, Nanded Pharmacy College, Nanded. Swami Ramanand Teerth Marathwada University, Nanded, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/aprj/2024/v12i5268>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc. are available here: <https://www.sdiarticle5.com/review-history/120902>

Review Article

Received: 07/06/2024

Accepted: 10/08/2024

Published: 19/08/2024

ABSTRACT

Barleria Prionitis Linn., commonly referred to as "Vajradanti," is a thorny shrub native to Southern Asia and certain African regions. This herb has a long history of use in traditional medicine systems, particularly Ayurveda, where it has been employed to address a wide array of health concerns

⁺⁺ Principal;

*Corresponding author: E-mail: fulchandk22699@gmail.com;

Cite as: Khandre, Vaishnavi V., Pallavi L. Bansod, Fulchand V. Kajale, N. B. Ghiware, and A. B. Roge. 2024. "A Review of *Barleria Prionitis* L. A Rarely Known Plant With Potential of Phytochemical and Pharmacological Assets". *Asian Plant Research Journal* 12 (5):39-47. <https://doi.org/10.9734/aprj/2024/v12i5268>.

including respiratory ailments, fever, liver disorders, and dermatological issues. A comprehensive review of scientific literature underscores the plant's rich phytochemical composition. It is a veritable treasure trove of bioactive compounds, such as tannins, saponins, flavonoids, glycosides, alkaloids, phenols, and specific compounds like β -sitosterol, lupeol, syringic acid, and vanillic acid. This chemical diversity is believed to underpin the herb's broad spectrum of pharmacological activities. Research has unveiled the potential of *Barleria Prionitis* in addressing a variety of health challenges. Studies have demonstrated its antimicrobial, antiparasitic, anti-inflammatory, antioxidant, and hepatoprotective properties. Furthermore, the herb has shown promise in managing metabolic disorders like diabetes, as well as gastrointestinal issues. Its analgesic and antiarthritic effects have also garnered attention. Despite the compelling evidence supporting the therapeutic efficacy of *Barleria Prionitis*, its full potential remains largely untapped. Further research is imperative to elucidate the underlying mechanisms of action, optimize extraction processes, and develop standardized formulations. Moreover, clinical trials are necessary to validate the traditional claims and establish the herb's safety and efficacy in modern healthcare. By unlocking the secrets of *Barleria Prionitis*, researchers and healthcare providers can potentially develop innovative and sustainable therapeutic solutions for a range of diseases.

Keywords: *Barleria prionitis*; *Acanthaceae*; *Antidiabetic*; *Anthelmintic*; *Antifertility*; *Antiviral*.

1. INTRODUCTION

Plants have served as both sustenance and medicine for millennia, with a vast array of species offering potential therapeutic benefits. India, a global biodiversity hotspot, is home to an abundance of medicinal plants. The contemporary resurgence of traditional medicine, driven by concerns over synthetic drug adverse effects, has ignited renewed interest in natural products. *Barleria Prionitis* L., a widely distributed Asian and African plant, holds a significant place in traditional medicine systems for treating various ailments [1]. Despite its historical use, comprehensive scientific research into its pharmacological and phytochemical properties remains relatively scarce. (Portal, India Biodiversity, 2020). This study aims to comprehensively elucidate the therapeutic potential of *Barleria Prionitis* L. by conducting a thorough investigation of its phytochemical constituents and pharmacological activities [2-5]. Specific objectives of this research encompass a detailed phytochemical profiling of the plant, evaluation of its pharmacological properties across multiple biological assays, and establishment of correlations between its phytochemical composition and observed biological effects. [6] By bridging this knowledge gap, this study contributes to the scientific validation of *Barleria Prionitis*' traditional medicinal applications and its potential as a promising source of novel therapeutic agents [7-12]. This comprehensive approach will not only enhance our understanding of the plant's pharmacological profile but also lay the groundwork for future studies focused on

isolating and characterizing bioactive compounds, optimizing extraction processes, and developing standardized formulations for clinical applications [13-17]. Ultimately, this research aspires to contribute to the development of safe, effective, and sustainable herbal-based therapies.

2. AIMS AND OBJECTIVES

To carried out the medicinal, pharmacological, and phytochemical properties of *Barleria* plant much possible. To explain the traditional uses of *Barleria. prionitis* in different medicinal systems. To provide a detailed botanic description of *Barleria. prionitis*. To discover the phytochemical constituents in *Barleria. prionitis*. To give more information about literature of this plant.

Synonyms: [18].

Table 1. Synonyms

Bengali	Kantajinti, Peetjhanti
English	Yellow nail, dye plant, Porcupine flower
Gujarati	Kantashila
Kannada	Karunta, Mullugorante
Malayalam	Chemmulli, Varelmutti
Marathi	Kate koranti, Kholeta, Koranta, Pivala koranta
Telugu	Mullugorintachettu

2.1 The Macroscopic Characters of *Barleria Prionitis* L

Barleria Prionitis L. is a shrub typically reaching a height of approximately 1.5 meters. The plant's

stems are cylindrical, exhibiting a rigid structure and a smooth texture. Their coloration varies from green to light tan. The leaves are elliptical to ovate in shape, with dimensions ranging from 3 to 10 centimetres in length and 1.5 to 4 centimetres in width [19]. A notable feature is the presence of 5 to 20 millimetre long, pale spines located in the axils of the leaves. The inflorescence is a terminal cluster of tubular flowers, each measuring around 3 to 4

centimetres in length [20]. These flowers are typically vibrant yellow or white, pink in colour. Following pollination, the plant develops ovoid capsules, approximately 8 millimetres long and 5 millimetres wide. These capsules are characterized by a covering of matted hairs and culminate in a sharp, pointed tip. [21].

Taxonomical Classification: [22].



Image. 1. *Barleria Prionitis* L.

Table 2. The macroscopic characters of *Barleria prionitis* Linn. Leaf

Sr.No.	Macroscopic Parameter	Observation
1	Colour	Dark green on upper surface and pale green on inferior surface
2	Odor	Characteristic
3	Taste	Tasteless
4	Habit	Erect; Perennial shrub
5	Shape	Ovate elliptical
6	Dimensions	3-14 × 1-3.5 cm
7	Apex	Acute or acuminate
8	Margin	Entire
9	Venation	Reticulate
10	Leaf base	Cuneate or Attenuate
11	Petiole	Petiolate

Table 3. Taxonomical classification

01.	Kingdom	Plantae
02.	Subkingdom	Tracheobionta
03.	Division	Magnoliophyta

04.	Class	Magnoliopsida
05.	Subclass	Asteridae
06.	Order	Scrophulariales
07.	Family	Acanthaceae
08.	Genus	<i>Barleria</i> Species - <i>Prionitis</i> .
09.	Botanical name	<i>Barleria prionitis</i> Linn
10.	Family	Acanthaceae (Acanthus family)
11.	Synonyms	<i>Barleria appressa</i> , <i>Barleria coriacea</i> ,

3. MATERIALS AND METHODS

3.1 Phytochemical Profile of *Barleria Prionitis* L

Barleria Prionitis L. has emerged as a promising phytotherapeutic agent due to its rich reservoir of secondary metabolites. The aerial parts of the plant, particularly the flowering tops and leaves, have been the primary focus of phytochemical investigations [23]. Potassium salts constitute a significant component of the plant's elemental composition, contributing to its overall therapeutic profile. Beyond these primary metabolites, a diverse array of secondary metabolites has been identified through a combination of traditional phytochemical screening methods and advanced analytical techniques. Flavonoids, alkaloids, tannins, saponins, terpenoids, steroids, essential oils, and phenolic compounds are the major classes of secondary metabolites reported in *Barleria Prionitis* L. [24]. These compounds have been qualitatively detected using a repertoire of colorimetric and precipitation reactions.

3.2 Test for the Detection of Phytochemical:

3.2.1 Test for alkaloids

- Wagner's Reagent: This reagent is a solution of iodine in potassium iodide. A reddish-brown precipitate indicates a positive test for alkaloids.
- Mayer's Reagent: This is a solution of potassium tetraiodomercurate (II). A cream-colored precipitate indicates a positive test for alkaloids.
- Dragendorff's Reagent: This reagent is a solution of bismuth subnitrate in potassium iodide and nitric acid. A reddish-orange precipitate indicates a positive test for alkaloids.

- Flavonoids by testing with alkaline reagent and Shinoda test.
- Saponins by Foam Test. A persistent foam that lasts for at least 10 minutes indicates the presence of saponins.
- di-nitro phenylhydrazine test for Terpenoids
- Phytosterol by Liebermann's test and Liebermann–Burchard test.
- Phenolic compounds and tannins by lead-acetate test, FeCl₃, and bromine water test.
- Essential oil by Sudan III test.
- Amino acids and proteins by Millon's test, Biuret test, and Ninhydrin test.
- Carbohydrates by Benedict's test, Fehling's test and Molisch test.
- Glycosides by Borntrager's test and legal's test.

4. RESULTS

4.1 Medicinal Uses

The flowers of this plant have garnered particular attention for their therapeutic properties. They are traditionally employed as a potent remedy for fever, a condition characterized by elevated body temperature. Additionally, the flowers have been used to manage respiratory ailments, which encompass a range of conditions affecting the airways, such as asthma, bronchitis, and the common cold. [25]. Furthermore, the anti-inflammatory properties of the flowers are believed to provide relief from joint pains, a common symptom associated with conditions like arthritis. The root of *Barleria Prionitis* Linn. has found specific application in oral health care. A mouthwash concocted from the root extract has been traditionally used to mitigate toothache and combat gum diseases, conditions characterized by inflammation and bleeding of the gum tissues [26,27]. Beyond the flowers and roots, the entire

Table 4. Activities of Phytoconstituents Isolated from *Barleria Prionitis* L

Plant Part	Phytoconstituent	Class	Activity	Ref.
Aerial part	Barlerinoside	Phenylethanoid Glycoside	Glutathione S-Transferase inhibitory activity	[28]
	Lupulinoside	Iridoid Diglucoside	Antioxidant activity	
	7-methoxydideroside	Secoiridoids	Antioxidant activity, antiviral activity	
	Balarenone	Terpenoid	Glutathione S-transferase and acetylcholinesterase inhibitory activity, antibacterial activity	[29]
	Lupeol	Triterpene	Anti-inflammatory and anti-cancer, glutathione s-transferase and acetylcholinesterase inhibitory activity, antibacterial activity	
	Apigenin 7-O- β -D-glucoside	Glycosyloxy Flavone	Antibacterial activity, anti-Inflammatory activity, ant-Oxidant activity	[29]
	Luteolin-7-O-glucoside	Flavone	Antibacterial activity, antioxidative activity, antimicrobial activity, hepatoprotective activity, anti-inflammatory activity	[28]
	Pipataline	Terpenoid	Enzyme inhibitory activity, antioxidant activity	[29]
	Barlerin	Iridoid Glycosides	Antioxidant activity, antiviral activity, anti-cancer activity, enzyme inhibitory activity, anti-inflammatory activity	[28]
	Acetyl Barlerin		Antioxidant activity, antiviral activity, anticancer activity, enzyme inhibitory activity, anti-inflammatory activity	[30]
	Shanzhiside methyl ester		Glutathione S-Transferase inhibitory activity, antioxidant activity	

plant, including the leaves, has been incorporated into traditional Indian medicine [8,31-33].

4.2 Chemical Constituent

The *Barleria Prionitis* leaves and flowering tops are rich source of potassium salt [34] Preliminary phytochemical analysis of hydro methanolic extract of *B. prionotis* whole plant indicated presence of flavonoids, glycosides, saponins, tannin and steroids [21]. Delving deeper into the plant's chemical composition, researchers have successfully isolated specific compounds, including balarenone, pipataline, lupeol, prioniside A, and prioniside B. These molecules represent promising targets for further investigation and potential drug development. Additionally, chromatographic studies have unveiled the presence of iridinoid glycosides, such as acetyl barlerin, expanding the repertoire

of bioactive compounds within this species. Chromatography study of alcohol extract of *Barleria prionitis* revealed the presence of iridinoid glycosides such as acetyl barlerin [26,27].

4.3 Traditional Uses

Several traditional uses of this plant are as follows:

- **Disorders of respiratory system:** [26,27]. Dried bark as well as roots are used as expectorant to treat whooping cough, Ash of the whole plant is used for asthma.
- **Pain fever and inflammation:** [29] Paste of the seeds is used to treat oedema, entire plant is used to treat tonsillitis, juice of the leaves is used to treat fever.

4.4 Pharmacological Activities of *Barleria prionitis* Linn.

1. **Anti-inflammatory and anti-nociceptive activity:** [26,27]. 50% ethanolic extract of the flower of *B. prionitis* in experimental animals shows anti-inflammatory as well as anti-nociceptive activity.
2. **Mast Cell Stabilization and Membrane Protection Activity:** [34]. Whole plant hydro-alcoholic extract of *Barleria prionitis* L. shows mast cell stabilization and membrane protection activity.
3. **Anti-bacterial Activity:** [25]. The methanolic extract from fresh callus cultures of *Barleria prionitis* which was cultured on MS medium fortified with IAA (2.0 mg/l) and BAP (1.5 mg/l) for 2 - 12 weeks was observed and examined for the potency of their phytochemicals against dental caries pathogens. It kills dental caries producing pathogens and shows anti-bacterial activity.
4. **Anticataract Activity:** [24]. Oral administration of *Barleria prionitis* significantly delayed the onset and progression of cataract in selenite as well as galactose induced cataract in pups.
5. **Antifertility Studies:** [35]. Oral administration of root extract of *Barleria prionitis* L. to male rats (100 mg/rat per day) for the period of 60 days did not cause body weight loss. The root extract brought about an interference with spermatogenesis. The round spermatids were decreased by 73.6% (P50.001). No significant change was found in the population of secondary spermatocytes.
6. **Antifungal Activity:** Pal, Anita, et al. [36]. Natural dye extracted from aerial parts of *Barleria prionitis* and different kinds of textile fabrics dyed with the natural dye were investigated for their antifungal activity.
7. **Antimicrobial Activity:** Panchal, Priyanka et al. [37]. Whole plant of *Barleria prionitis* L. whole plant shows anti-bacterial property in ethanolic extract.

8. **Antioxidant Activity:** [38]. Different extract of *Barleria prionitis* L. leaves and stem shows anti-oxidant activity.

9. **Central Nervous System Activity:**[39]. The 70% ethanol extract of leaves of *Barleria prionitis* Linn (Acanthaceae) in Swiss albino mice. General behaviour was studied using actophotometer. According to the study, it was observed that the test drug has the stimulant activity.

5. DISCUSSION

The present review underscores the historical and traditional significance of *Barleria prionitis* in addressing a diverse array of health concerns, notably in the Indian medicinal repertoire. The plant's efficacy in managing ailments such as toothache, wounds, and joint pain provides a compelling rationale for contemporary scientific investigation. The phytochemical profile of *B. prionitis*, characterized by the presence of glycosides, saponins, flavonoids, steroids, and tannins, offers a promising foundation for elucidating the plant's pharmacological properties. These bioactive compounds are well-documented for their therapeutic potential in various pharmacological activities, including anti-inflammatory, antioxidant, and antimicrobial effects. Future research should focus on isolating and characterizing the bioactive compounds, conducting pharmacological evaluations, and exploring potential therapeutic applications. By bridging the gap between traditional knowledge and modern scientific inquiry, the full potential of *B. prionitis* can be realized.

6. CONCLUSION

The review of *Barleria prionitis* offers valuable insights into the plant's potential as a therapeutic agent. While further research is needed to fully understand its medicinal potential, the conclusions presented in the review provide a strong foundation for future investigations. A multidisciplinary approach involving ethnobotany, phytochemistry, pharmacology, and clinical research is essential to reveal the full therapeutic promise of this underutilized plant. The article is informative and resourceful. It will definitely help the practitioners and researchers to do more works in the field. This study is important in terms of scientific research because of the side effects of modern synthetic drugs.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Vasoya U. (2012). Pharmacognostical and physicochemical studies on the leaves of *Barleria prionitis* (L.). International Journal of Pharmaceutical Sciences and Research. 2012;3(7):2291.
- Banerjee D, Maji AK, Mahapatra S, Banerji P. *Barleria prionitis* Linn.: A review of its traditional uses, phytochemistry, pharmacology and toxicity. Research Journal of Phytochemistry. 2012; 6(2):31-41.
- Ahmad S, Rageeb M, Usman M, Jain BV, Shaikh MZ. *Barleria Prionitis*: It's pharmacognosy, phytochemicals and its potential beneficial effects in common oro-dental diseases. Journal of Current Pharma Research. 2000;11(1): 3941-3966.
- Amarasiri SS, Attanayake AP, Arawwawala LD, Mudduwa LK, Jayatilaka KA. *Barleria prionitis* L. extracts ameliorate doxorubicin-induced acute kidney injury via modulation of oxidative stress, inflammation, and apoptosis. Journal of Traditional and Complementary Medicine. 2023;3(5): 500-510.
- Amarasiri SS, Attanayake AP, Mudduwa LK, Jayatilaka KA. Acute and sub-acute toxicological evaluation of *Barleria prionitis* Linn. in Wistar albino rats. Ceylon Journal of Science. 2023;52(4): 445-454.
- Shukla S, Gunjegaokar SM. Pharmacognostical and pharmacological profiling of *Barleria prionitis* Linn. Journal of Biological Sciences and Medicine. 2018;4(1):41-50.
- Dheer R, Swarnkar SK, Syeed F. Chromatographic analysis of *Barleria prionitis* Linn. Research Journal of Pharmacy and Technology. 2019;12(8): 3679-3686.
- Ata A, Kalhari KS, Samarasekera R. Chemical constituents of *Barleria prionitis* and their enzyme inhibitory and free radical scavenging activities. Phytochemistry Letters. 2009;2(1): 37-40.
- Banerjee S, Banerjee S, Jha GK, Bose S. *Barleria prionitis* L.: An illustrative traditional, phytochemical and pharmacological review. The Natural Products Journal. 2021;11(3): 258-274.
- Choudhary M, Kumar V, Gupta PK, Singh S. Anti-arthritis activity of *Barleria prionitis* Linn. leaves in acute and chronic models in Sprague Dawley rats. Bulletin of Faculty of Pharmacy, Cairo University. 2014;52(2): 199-209.
- Diwan PD, Gadhikar YA. Assessment of phytochemical composition and antibacterial activity of different extracts of *Barleria prionitis* leaves against oral microflora to improve dental hygiene. Asian J. Pharm. Clin. Res. 2012;5(2):182-184.
- El-Emary NA, Makboul MA., Abdel-Hafiz MA, Ahmed AS. Phytochemical study of *Barleria cristata* L. and *Barleria prionitis* L. cultivated in Egypt. Bulletin of Pharmaceutical Sciences. Assiut. 1990;13(1):65-72.
- Farrukh S. A review about *Barleria Prionitis*: A rare known shrub with potential medicinal properties. J Pharm Sci Drug Discov. 2022;1(1):1-7.
- Gangaram S, Naidoo Y, Dewir YH, El-Hendawy S. Phytochemicals and biological activities of *Barleria* (Acanthaceae). Plants. 2021;11(1):82.
- India BP. Online Portal of India Biodiversity. Online Portal of India Biodiversity; 2020.
- Kosmulalage KS, Zahid S, Udenigwe CC, Akhtar S, Ata A, Samarasekera R. Glutathione S-transferase, acetylcholinesterase inhibitory and antibacterial activities of chemical constituents of *Barleria prionitis*. Zeitschrift für Naturforschung B. 2007;62(4): 580-586.
- Naryan S, Nikam, P. A review on, pharmacological activities of *Barleria prionitis* Linn. World Journal of

- Pharmaceutical Research. 2018; 7:166-180.
18. Singh MP, Panda H. Medicinal herbs with their formulations. Daya Books; 2005.
19. Khare CP. Indian medicinal plants: an illustrated dictionary. Springer Science & Business Media. 2008;699-700.
20. Kapoor A, Shukla S, Kaur R, Kumar R, Lehra KS, Kapoor S. Preliminary phytochemical screening and antioxidant activity of whole plant of *Barleria prionitis* Linn. Int. J. Adv. Pharm. Biol. Chem. 2014;3(2):410-419.
21. Sharma P, Shrivastava B, Sharma G, Jadhav H. Phytochemical and ethnomedical values of *Barleria prionitis* L: An overview. J Harmon Res Pharm. 2013; 2:190-93.
22. Bhogaonkar PY, Lande SK. Anatomical characterization of *Barleria prionitis* Linn.: A well-known medicinal herb. In Biol. Forum Int. J. 2012; 4:1-5.
23. Ghatapanadi SR, Johnson N, Rajasab AH. Documentation of folk knowledge on medicinal plants of Gulbarga district, Karnataka; 2011.
24. Atif, Mohammed, Rahman SA, Ahmed MI, Mahmood SB, Azharuddin Mohammed. Anticataract potential of *Barleria prionitis*: In vivo study. Int J Pharm Pharm Sci. 2015;7(2):100-55.
25. Kumari Purnima, Yadav Poonam, Arya Arvind, Kumar, Sandeep. *In vitro* callus production and anti-bacterial activity of *Barleria prionitis* Linn. against dental caries pathogens. Int J Bot Res. 2013;3(4): 1-6.
26. Jaiswal SK, Dubey MK, Das S, Verma AR, Vijayakumar M, RAO CV. Inflammatory and Anti Nociceptive Activity. International Journal of Pharma and Bio Sciences. 2010; 1:2.
27. Jaiswal SK, Dubey MK, Verma AK, Das S, Vijaykumar M, Rao CV. Evaluation of iridoid glycosides from leaves of *Barleria prionitis* as an anti-diarrhoeal activity: An ethnopharmacological study. Int. J. Pharm. Sci. 2010;2(3): 680-686.
28. Ata A, Naz S, Elias EM. Naturally occurring enzyme inhibitors and their pharmaceutical applications. Pure and Applied Chemistry. 2011;83(9): 1741-1749.
29. Panchal PK, Sharma N, Singh K. Antibacterial effects of aerial part of *barleria prionitis* with ethanol extarct on pathogenic strains. World J Pharm Pharm Sci. 2017; 6:1147-55.
30. Pandey K, Gupta H, Kamble B. *Barleria prionitis*: Journey from Ayurveda to modern medicine. The Natural Products Journal. 2018;8(2):109-130.
31. Talele BD, Mahajan RT, Chopda MZ, Nemade NV. Nephroprotective plants: a review. Int J Pharm Pharm Sci, 2000;4(1), 8-16.
32. Chen JL, Blanc P, Stoddart CA, Bogan M, Rozhon EJ, Parkinson N, Kernan MR. New iridoids from the medicinal plant *Barleria prionitis* with potent activity against respiratory syncytial virus. Journal of Natural Products. 1998;61(10): 1295-1297.
33. Aneja KR, Joshi R, Sharma C. Potency of *Barleria prionitis* L. bark extracts against oral diseases causing strains of bacteria and fungi of clinical origin. New York Sci J. 2010;3(11):5-12.
34. Maji AK, Bhadra S, Mahapatra S, Banerji P, Banerjee D. Mast cell stabilization and membrane protection activity of *Barleria prionitis* L. *Pharmacognosy Journal*. 2011;3(24):67-71.
35. Gupta RS, Kumar P, Dixit VP, Dobhal MP. Antifertility studies of the root extract of the *Barleria prionitis* Linn in male albino rats with special reference to testicular cell population dynamics. Journal of Ethnopharmacology. 2000;70(2): 111-117.
36. Pal A, Kumar R, Upadhyay L, Tripathi YC. Antifungal activity of natural dye from aerial biomass of *Barleria prionitis* L. and dyed fabrics. Iran. J. Chem. Chem. Eng. Research Article. 2018;37(1).
37. Panchal Priyanka, Singh, Kamal. Antimicrobial activity of *Barleria prionitis* on pathogenic strains. Int. J. Curr. Pharm. Res. 2015;7(4):73-75.
38. Sharma P, Sharma GN, Shrivastava B, Jadhav HR. Evaluation of antioxidant potential of *Barleria prionitis* leaf and stem. American Journal of Phytomedicine and Clinical Therapeutics. 2014;2(10):1177-86.

39. Gangopadhyay A, Malakar J, Ghosh A, Deb J, Dey S, Datta S, Datta PK. The central nervous system activity of *Barleria prionitis* Linn. on the locomotor activity of swiss albino mice using actophotometer. Int J Pharm Biol Sci Arch. 2012;3(2): 403-405.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/120902>