



# **Effect of Corm and Soil Treatments with Organic Liquid Formulations on Performance of Elephant Foot Yam in Jharkhand, India**

**Saikat Biswas <sup>a\*</sup>, Rupa Das <sup>b</sup> and Raghava Thakur <sup>a</sup>**

<sup>a</sup> Faculty of Agriculture, Rural and Tribal Development, Ramakrishna Mission Vivekananda Educational and Research Institute, Morabadi, Ranchi, Jharkhand-834008, India.

<sup>b</sup> Department of Seed Science and Technology, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal-741252, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJPSS/2023/v35i52827

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97278>

**Original Research Article**

**Received: 05/01/2023**

**Accepted: 09/03/2023**

**Published: 13/03/2023**

## **ABSTRACT**

A field experiment was carried out at the farm of Divyayan Krishi Vigyan Kendra, Ramakrishna Mission Ashrama, Morabadi, Ranchi during kharif season of 2022 with 10 treatments (T<sub>1</sub>: control or no treatment, T<sub>2</sub>: Corm treatment with Beej Sanjeevani, T<sub>3</sub>: Corm treatment with Beejamrit, T<sub>4</sub>: Soil application of Jiwamrit, T<sub>5</sub>: Soil application of Panchagavya, T<sub>6</sub>: Soil application of Jiwamrit + Panchagavya, T<sub>7</sub>: Corm treatment with Beej Sanjeevani + soil application of Jiwamrit, T<sub>8</sub>: Corm treatment with Beej Sanjeevani + soil application of Panchagavya, T<sub>9</sub>: Corm treatment with Beejamrit + soil application of Jiwamrit, T<sub>10</sub>: Corm treatment with Beejamrit + soil application of Panchagavya) placed in three times replicated randomized block design to study the growth, yield and production economics of elephant foot yam under uses of organic corm and soil treatments. Results revealed that corm treatment with Beejamrit + soil application of Jiwamrit resulted in higher

\*Corresponding author: E-mail: sbsaikatbiswas27@gmail.com;

growth attributes, corm length (23.3 cm), diameter (26.9 cm) and yield (32.8 t/ha). This corm + soil treatment also recorded the highest gross return (INR.2,62,400/ha), net return (INR.1,69,189/ha) and B:C (2.82). It also increased soil microbial population dynamics especially total bacteria ( $69.7 \text{ CFU} \times 10^6/\text{g}$  of soil).

**Keywords:** Beej Sanjeevani; Beejamrit; Jiwarmit; economics; elephant foot yam; Panchagavya; yield.

## 1. INTRODUCTION

Long term exploitation of natural resources under intensive agriculture not only deteriorates agricultural productivity but also hampers the environment safety. Use of chemicals in crop production results in deterioration of soil health and overall environment due to toxic residues. Continuous and pointless application of chemical fertilizers causes impairments of soil physical, chemical and biological properties, loss of soil productivity and environmental pollution, health issues etc. Use of chemicals is beneficial for short term basis and harmful for long term sustainable agricultural production [1]. Khambalkar et al. [2] stated that the harmful impacts of chemical fertilizers in intensive agriculture such as depletion of reserve pool of carbon, secondary and micro-nutrients, resulting in degradation of soil productivity and fertility. Further, increase of market price of chemicals is another issue under energy crisis scenario. In India, land degradation is a common issue which is contributed to a major extent through faulty agricultural practices [3]. To ensure environment safety without loss of crop productivity eco-friendly alternatives of chemical farming like organic farming, natural farming, integrated nutrient management, biodynamic farming etc. show great prospects in current agriculture scenario [4]. Addition of organic formulations or proper utilization of natural resources can boost up the growth and productivity by reviving soil productivity [5].

The concept of natural farming is receiving spotlight in recent days, which encourages use of liquid formulations for corm and treatment, mulching (Acchadana), and soil moisture conservation (Whapasa) [4,6]. Various liquid organic formulations like Panchagavya, Sasyagavya, Kunapajala, and Jiwarmit as soil/foliar use and Beejamrit and Beej Sanjeevani as seed treatment are showing good potential for improvement of microbial activity and population, soil physical and fertility elevation, which finally reflect on crop growth, yield and quality [7]. They contain various beneficial micro-organisms, macro and micro-nutrients, plant growth

promoters, antioxidant and plant protection properties etc [8]. These liquid manures are prepared from on-farm/local raw materials or products like cow dung, cow urine, cow ghee, cow milk, cow curd, crop field and livestock wastes and some other low-cost materials through fermentation and/or decomposition. Application of these organic formulations leads to healthy plant-microbe interactions [9,10] and thus can reduce or omit the use of chemicals [7].

Elephant foot yam (EFY), [*Amorphophallus paeoniifolius* (Dennst.) Nicolson] is one of the major tropic aroids grown in India since long specially in Andhra Pradesh, Tamil Nadu, West Bengal and Kerala with productivity potential of 30-100 t/ha [11]. The tubers, leaves and tender pseudo stem are used as vegetables. The tuber is cheap source of starch, calcium, phosphorus and vitamin A, protein etc and can substitute potato to a great extent specially in economically poor areas. The growth and yield of EFY largely dependent on nutrient management and water availability. Proper corm and soil treatments using different organic liquid formulations can play important role in booting the growth and yield of EFY. With this view, the present experiment was placed at the farm of Divyayan Krishi Vigyan Kendra, Ramakrishna Mission Ashrama, Morabadi, Ranchi during kharif season of 2022 to identify the most suitable corm and soil treatments for growth and productivity of EFY.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Details

The field experiment was conducted at the farm of Divyayan Krishi Vigyan Kendra, Ramakrishna Mission Ashrama, Morabadi, Ranchi during kharif season of 2022 to evaluate the response of elephant foot yam to corm + soil treatments with various organic liquid formulations under 13 years old organic field. The experimental soil was clay loam textured, red and lateritic in nature having 6.70 of pH, 1.23% organic carbon, 396.14 kg/ha of available N, 147.2 kg/ha of available  $\text{P}_2\text{O}_5$  and 245.71 kg/ha available  $\text{K}_2\text{O}$ . The

experiment was placed in randomized block design with 10 treatments ( $T_1$ : control or no treatment,  $T_2$ : Corm treatment with Beej Sanjeevani,  $T_3$ : Corm treatment with Beejamrit,  $T_4$ : Soil application of Jiwamrit,  $T_5$ : Soil application of Panchagavya,  $T_6$ : Soil application of Jiwamrit + Panchagavya,  $T_7$ : Corm treatment with Beej Sanjeevani + soil application of Jiwamrit,  $T_8$ : Corm treatment with Beej Sanjeevani + soil application of Panchagavya,  $T_9$ : Corm treatment with Beejamrit + soil application of Jiwamrit,  $T_{10}$ : Corm treatment with Beejamrit + soil application of Panchagavya), replicated three times. The preparation and application of different organic liquid formulations was shown in Table 1. The EFY variety 'Gajendra' was planted on individual plot size was 4 m x 3 m using ( $\geq 500$  g weight whole/cut corm having bud) on 2<sup>nd</sup> May, 2022 at a spacing of 75 cm x 75 cm and harvested on Nov 09, 2022.

Prior to planting, 20 t/ha FYM was applied during land preparation. Corm (corm) treatments with two organic liquid formulations were made for 30 minutes before planting (Fig 1a). Soil application of organic liquid formulations were made at 15 days interval up to 5 months after planting (MAP). Weeding was periodically done up to 3 MAP. Irrigation (4 cm depth) was given for 4 times during initial 3 months at an interval of 20 days along with leaf and paddy straw mulching to conserve and provide moisture for

crop growth during pre-monsoon period. During rainy season, the crop utilized rainwater for its growth. Other agronomic and plant protection measures were taken as per the standard organic farming practices recommended for this region.

## 2.2 Observations Recorded

Observations on plant population (per plot) was taken at harvest. Different growth attributes like plant height (cm), canopy width (cm) and pseudo stem girth at collar region (cm) were taken using measuring tape/scale at 3 MAP and at harvest. For recording of observations, five random plants were selected from each plot and tagged (Fig 1b). Observations were taken from those five plants. Corm diameter and length as well as corm yield were recorded at harvest. Production economics of EFY consisted of cost of cultivation, gross return, net return and benefit: cost (B:C) which were computed based on following formulae:

- Cost of cultivation (INR./ha) = Cost involved in purchase of inputs as well as practices
- Gross return (INR. /ha) = Yield (kg/ha) x Market price (INR./kg)
- Net return (INP. /ha) = Gross return (INR./ha) - Cost of cultivation (INR./ha)
- B:C = Gross return (INR./ha) / Cost of cultivation (INR./ha)



Fig. 1a. Corm treatment of elephant foot yam




Fig. 1b. Elephant foot yam plants

**Table 1. Preparation of various organic liquid formulations**

Organic liquid formulation	Preparation	Application
Beej Sanjeevani	200 g cow dung, 200 ml cow urine and 800 ml water are mixed together and stirred twice a day (morning and evening), clockwise and anti-clockwise for 7 days.	It was used as corm treatment @ 5% by diluting the mother solution with water.
Beejamrit	200 g cow dung, 200 ml cow urine, 100 g lime, one pinch of live soil and 800 ml water are mixed together. It was followed by stirring twice a day (morning and evening), clockwise and anti-clockwise for 7 days.	It was used as corm treatment @ 5% by diluting the mother solution with water.
Jiwamrit	10 kg fresh cow dung, 10 litres of cow urine, 2 kg jaggery, 2 kg pulse flour, a fist of organically cultivated soil and 200 litres of water are taken and mixed thoroughly in a container. This mixture is kept in shade for fermentation for 48 hours with regular stirring twice a day (morning and evening), clockwise and anti-clockwise.	It was applied to soil @ 10% by diluting the mother solution with water.
Panchagavya	Five cow-based ingredients <i>i.e.</i> cow dung, cow urine, milk, curd and ghee were added in a plastic container @ 5:3:2:2:1 ratio and incubated for 7-9 days. The mixture was stirred with a stick every day two times during morning and evening, clockwise and anti-clockwise.	It was applied to soil @ 10% by diluting the mother solution with water.

Soil microbial population (total bacteria and fungi) dynamics during initial (before experiment) and after harvest of crop were estimated at the laboratory of Ramakrishna Mission Vivekananda Educational and Research Institute, Morabadi, Ranchi, Jharkhand. The population of soil rhizospheric micro-organisms (bacteria and fungi) were analysed through counting on agar plates as number of viable cells per gram of soil using Thronton's agar medium and Martin-Rose Bengal Streptomycin agar medium respectively using serial dilution and pour plate process [12] followed by incubation at  $28\pm 1^\circ\text{C}$ . The populations were counted at 5<sup>th</sup> day of incubation.

### 2.3 Statistical Analysis

The data were subjected to statistical analysis using ANOVA method [13] and treatment means were compared using critical difference at 5% level of significance.

## 3. RESULTS AND DISCUSSION

### 3.1 Plant Population and Growth Attributes

Experimental results (Table 2) expressed that there existed non-significant variations in plant

population of EFY at harvest among different organic corm + soil treatments and control with highest plant population observed from corm treatment with Beejamrit + soil application of Jiwamrit ( $T_9$ ) (21.3/plot) and the lowest was from control ( $T_1$ ) (18.6/plot). However, different growth attributes of EFY at 3 MAP and at harvest showed significant variations among different organic corm + soil treatments and control. Elephant foot yam showed variation in growth attributes as per the microbial and nutritional properties of the organic formulations used. Corm treatment with Beejamrit ensured high plant population which might be due to its high nutrients, growth promoters, and beneficial microorganisms possibly resulting in enhancement of metabolic activities like water intrusion to planting material and breakdown of storage materials of corms through enzymes like lipase, proteinase, phosphatase, and hydrolase [14]. Improved metabolic activity led to better emergence of plants. Further, presence of micro-organisms in organic formulations especially Beejamrit and Jiwamrit can protect crops against diseases during the early growth period [15]. Corm treatment with Beejamrit + soil application of Jiwamrit ( $T_9$ ) recorded the maximum plant height, canopy width and pseudo stem girth at collar region, being closely followed by corm treatment with Beejamrit + soil application of

Panchagavya ( $T_{10}$ ) which remained statistically similar to corm treatment with Beej Sanjeevani + soil application of Panchagavya ( $T_8$ ), corm treatment with Beej Sanjeevani + soil application of Jiwamrit ( $T_7$ ), soil application of Jiwamrit + Panchagavya ( $T_6$ ) and soil application of Jiwamrit ( $T_4$ ) in terms of plant height and canopy width at 3 MAP and at harvest. Lowest plant height, canopy width and pseudo stem girth at collar region were noted from control ( $T_1$ ). The presence of growth-regulating hormones and enzymes in Jiwamrit possibly facilitated the uptake of nutrients and moisture by plants leading to meristematic cell division and elongation, which was finally reflected to plant height [16]. Corm treatment with Beejamrit + soil application of Jiwamrit ( $T_9$ ) outperformed others as both the formulations were possibly rich in macro- and micro-nutrients, vitamins, amino acids, growth regulators and served as storage media for the beneficial microorganisms [17,18]. In soil, they perhaps positively influenced soil microbial activity for sustainable supply of nutrients to EFY [19]. Greater N availability and uptake might trigger the leaf chlorophyll synthesis for capturing more sunlight and  $CO_2$  resulting in high photosynthesis, dry matter production and good vegetative growth [20].

### 3.2 Yield Attributes and Yield

Yield attributes such as corm length and diameter and corm yield were taken at harvest and they showed significant variations among different corm + soil treatments (Table 3). It was found that different soil and corm treatments alone or in combination outperformed control in attaining yield attributes and yield of EFY. Among different organic formulations, Corm treatment with Beejamrit + soil application of Jiwamrit ( $T_9$ ) recorded the highest corm length (23.3 cm) and diameter (26.9 cm), closely followed by corm treatment with Beejamrit + soil application of Panchagavya ( $T_{10}$ ) (corm length: 21.7 cm and diameter: 25.4 cm). Control showed lowest of those (corm length: 16.8 cm and diameter: 18.3 cm). Corm yield also followed similar trend where  $T_9$  registered the maximum yield (32.8 t/ha) which was 73.5% higher than the control (18.9 t/ha). It closely followed by  $T_{10}$  (31.1 t/ha). The sequence of treatments for expressing different yield attributes and yield was  $T_9 > T_{10} > T_7 > T_8 > T_4 > T_6 > T_5 > T_3 > T_2 > T_1$ .  $T_4$ ,  $T_6$ ,  $T_7$  and  $T_8$  remained statistically indifferent to each other. Yield attributes and yield were the direct reflection of crop growth. In the present study, corm treatment with Beejamrit might help in initial crop establishment, while soil application of

Jiwamrit as in case of  $T_9$  further helped in survivals, multiplications and activities of N-fixing, P-solubilising, cellulose decomposing and other beneficial microbes, which positively changed soil physical properties and chemical properties [21,4]. Soil application of Jiwamrit additionally might increase water-holding capacity of soil which helped the crop to utilize water during pre- and post-monsoon periods [6]. Along with these organic formulations, FYM was applied additionally at basal which might help in the early decomposition and quick release of nutrients at initial growth of the plants [22,23]. The adequate supply of nutrients over a longer period of time might not only facilitate the photosynthesis of the crop but also the translocation of photo-assimilates to economic parts *i.e.* corm [24,25].

### 3.3 Production Economics

Production economics of EFY (Table 4) indicated that use of organic liquid formulations either individually for corm and soil treatments or in combination incurred higher cost of cultivation of over control (INR.79,349/ha) because of the requirement of various raw materials for their preparations as well as applications. Among the treatments, corm treatment with Beejamrit + soil application of Panchagavya ( $T_{10}$ ) incurred maximum cost of cultivation (INR.96,487/ha) of EFY. It was perhaps due to the cost of various cow-based products (cow dung, cow ghee, cow urine, cow milk, cow curd) used for preparation of Panchagavya. However, corm treatment with Beejamrit + soil application of Jiwamrit ( $T_9$ ) recorded maximum gross return (INR.2,62,400/ha), net return (INR.1,69,189/ha) and B:C (2.82). The result was due to its positive and maximum influence on the growth and yield of EFY [26] and its efficacy for producing a greater return from corm yield than the expenses required [27,28]. The above treatment combination was closely followed by corm treatment with Beejamrit + soil application of Panchagavya ( $T_{10}$ ) (gross return: INR.2,48,800/ha, net return: INR.1,52,313/ha and B:C of 2.58). Control ( $T_1$ ), on the other hand, recorded lowest gross return (INR.1,51,200/ha), net return (INR.71,851/ha) and B:C (1.91) because of poor growth and yield of EFY.

### 3.4 Microbial Population Dynamics

The changes in total soil bacteria and fungi population before and after EFY cultivation were shown in Table 5. It was observed that except control, population of these microorganisms

enhanced at harvest as compared to initial (pre-experiment) status under uses of various organic liquid formulations as corm and soil treatments alone or in combination. In case of control, those declined after crop harvest. Among different organic liquid formulations, maximum increase of total bacteria (20.59%) and fungi (13.89%) over control (total bacteria:  $57.8 \text{ CFU} \times 10^6/\text{g}$  of soil; total fungi:  $21.6 \text{ CFU} \times 10^4/\text{g}$  of soil) were obtained from corm treatment with Beejamrit + soil application of Jiwamrit ( $T_9$ ) (total bacteria:  $69.7 \text{ CFU} \times 10^6/\text{g}$  of soil; total fungi:  $24.9 \text{ CFU} \times$

$10^4/\text{g}$  of soil). It was closely followed by corm treatment with Beejamrit + soil application of Panchagavya ( $T_{10}$ ) (total bacteria:  $67.5 \text{ CFU} \times 10^6/\text{g}$  of soil; total fungi:  $24.1 \text{ CFU} \times 10^4/\text{g}$  of soil). It might be due to positive influence of organic formulations on soil microbial growth. These formulations perhaps showed synergies with soil micro-organisms due to presence of high nutrients, beneficial micro-organisms, growth regulators like indole acetic acid, gibberellic acid, cytokinin, etc., which was earlier confirmed by many researchers [7,29,30].

**Table 2. Effect of corm and soil treatments with organic liquid formulations on growth of elephant foot yam**

Treatments	Plant population (/plot)	Plant height (cm)		Canopy width (cm)		Pseudo stem girth at collar region (cm)	
		3 MAP	At harvest	3 MAP	At harvest	3 MAP	At harvest
Control ( $T_1$ )	18.6	54.3	89.6	49.5	87.3	11.8	17.2
Corm treatment with Beej Sanjeevani ( $T_2$ )	20.3	62.1	98.9	59.3	102.1	13.1	19.8
Corm treatment with Beejamrit ( $T_3$ )	20.9	64.6	106.2	63.4	108.9	13.4	20.2
Soil application of Jiwamrit ( $T_4$ )	19.6	72.5	112.9	72.1	114.2	14.6	21.3
Soil application of Panchagavya ( $T_5$ )	19.3	69.1	108.7	69.1	110.5	13.8	20.5
Soil application of Jiwamrit + Panchagavya ( $T_6$ )	20.3	71.8	110.9	70.4	111.2	14.2	20.9
Corm treatment with Beej Sanjeevani + soil application of Jiwamrit ( $T_7$ )	20.9	74.3	116.8	72.8	118.3	15.8	22.7
Corm treatment with Beej Sanjeevani + soil application of Panchagavya ( $T_8$ )	21.0	73.3	114.3	72.6	115.7	15.1	22.3
Corm treatment with Beejamrit + soil application of Jiwamrit ( $T_9$ )	21.3	82.4	128.9	77.9	131.4	17.9	24.7
Corm treatment with Beejamrit + soil application of Panchagavya ( $T_{10}$ )	21.0	74.6	119.3	73.2	121.2	16.3	23.5
S.Em $\pm$	1.2	2.23	2.73	1.30	3.20	0.27	0.37
C.D. ( $p=0.05$ )	NS	6.7	8.2	3.9	9.6	0.8	1.1

**Table 3. Effect of corm and soil treatments with organic liquid formulations on yield attributes and yield of elephant foot yam**

Treatments	Corm length (cm)	Corm diameter (cm)	Corm yield (t/ha)
Control ( $T_1$ )	16.8	18.3	18.9
Corm treatment with Beej Sanjeevani ( $T_2$ )	18.2	20.4	23.1
Corm treatment with Beejamrit ( $T_3$ )	18.9	20.9	24.6
Soil application of Jiwamrit ( $T_4$ )	20.4	23.3	26.8
Soil application of Panchagavya ( $T_5$ )	19.3	21.8	25.7
Soil application of Jiwamrit + Panchagavya ( $T_6$ )	20.1	23.2	26.3
Corm treatment with Beej Sanjeevani + soil application of Jiwamrit ( $T_7$ )	21.4	24.9	29.5



Treatments	Corm length (cm)	Corm diameter (cm)	Corm yield (t/ha)
Corm treatment with Beej Sanjeevani + soil application of Panchagavya (T <sub>8</sub> )	20.6	23.5	27.1
Corm treatment with Beejamrit + soil application of Jiwamrit (T <sub>9</sub> )	23.3	26.9	32.8
Corm treatment with Beejamrit + soil application of Panchagavya (T <sub>10</sub> )	21.7	25.4	31.1
S.Em ±	0.43	0.57	1.23
C.D. (p= 0.05)	1.3	1.7	3.8

**Table 4. Effect of corm and soil treatments with organic liquid formulations on production economics of elephant foot yam**

Treatments	Cost of cultivation (INR./ha)	Gross return* (INR./ha)	Net return (INR./ha)	B:C
Control (T <sub>1</sub> )	79,349	1,51,200	71,851	1.91
Corm treatment with Beej Sanjeevani (T <sub>2</sub> )	84,456	1,84,800	1,00,344	2.19
Corm treatment with Beejamrit (T <sub>3</sub> )	84,762	1,96,800	1,12,038	2.32
Soil application of Jiwamrit (T <sub>4</sub> )	87,890	2,14,400	1,26,510	2.44
Soil application of Panchagavya (T <sub>5</sub> )	91,346	2,05,600	1,14,254	2.25
Soil application of Jiwamrit + Panchagavya (T <sub>6</sub> )	90,234	2,10,400	1,20,166	2.33
Corm treatment with Beej Sanjeevani + soil application of Jiwamrit (T <sub>7</sub> )	92,345	2,36,000	1,43,655	2.56
Corm treatment with Beej Sanjeevani + soil application of Panchagavya (T <sub>8</sub> )	95,327	2,16,800	1,21,473	2.27
Corm treatment with Beejamrit + soil application of Jiwamrit (T <sub>9</sub> )	93,211	2,62,400	1,69,189	2.82
Corm treatment with Beejamrit + soil application of Panchagavya (T <sub>10</sub> )	96,487	2,48,800	1,52,313	2.58

\*Market price of EFY: INR. 8.0/kg

**Table 5. Effect of corm and soil treatments with organic liquid formulations on soil microbial population dynamics of elephant foot yam field**

Treatments	Total bacteria (CFU × 10 <sup>6</sup> /g of soil)		Total fungi (CFU × 10 <sup>4</sup> /g of soil)	
	Initial	Harvest	Initial	Harvest
Control (T <sub>1</sub> )	57.8	53.4	21.6	19.1
Corm treatment with Beej Sanjeevani (T <sub>2</sub> )	57.8	64.7	21.6	22.1
Corm treatment with Beejamrit (T <sub>3</sub> )	57.8	65.2	21.6	22.5
Soil application of Jiwamrit (T <sub>4</sub> )	57.8	66.3	21.6	23.2
Soil application of Panchagavya (T <sub>5</sub> )	57.8		21.6	22.6
Soil application of Jiwamrit + Panchagavya (T <sub>6</sub> )	57.8	66.2	21.6	22.8
Corm treatment with Beej Sanjeevani + soil application of Jiwamrit (T <sub>7</sub> )	57.8	67.1	21.6	23.9
Corm treatment with Beej Sanjeevani + soil application of Panchagavya (T <sub>8</sub> )	57.8	67.0	21.6	23.6
Corm treatment with Beejamrit + soil application of Jiwamrit (T <sub>9</sub> )	57.8	69.7	21.6	24.9
Corm treatment with Beejamrit + soil application of Panchagavya (T <sub>10</sub> )	57.8	67.5	21.6	24.1
S.Em ±	-	0.57	-	0.23
C.D. (p= 0.05)	-	1.7	-	0.7

## 4. CONCLUSION

In the above study, organic liquid formulations as corm and soil treatments either alone or in combination proved their efficacies in influencing growth, yield, economics and soil residual microbial population of elephant foot yam cultivation. Based on the result, it is concluded that elephant foot yam can be cultivated treating corm with Beejamrit prior to planting and applying Jiwarmit @ 10% to soil at 15 days interval up to 5 months after planting for realizing best growth, corm yield, economic profitability and soil biological property.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Biswas S, Jana K, Agrawal RK, Puste AM. Effect of integrated nutrient management on growth attributing characters of crops under various oat-lathyrus intercropping system. The Pharma Innovation Journal. 2019;8(9):368-373.
2. Khambalkar PA, Tomar PS, Verma SK. Long term effects of integrated nutrient management on productivity and soil fertility in pearl millet (*Pennisetum glaucum*)-mustard (*Brassica juncea*) cropping sequence. Indian Journal of Agronomy. 2012;57(3):222–228.
3. Bhattacharyya R, Ghosh BN, Mishra PK, Mandal B, Rao CS, Sarkar D, Das K, Anil KS, Lalitha M, Hati KM, Franzluebbers AJ. Soil degradation in India: Challenges and potential solutions. Sustainability. 2015;7(4):3528-3570.
4. Liao J, Xu Q, Xu H, Huang D. Natural farming improves soil quality and alters microbial diversity in a cabbage field in Japan. Sustainability. 2019;3131(11):1-16.
5. Santhi R, Selvakumari G. Use of organic sources of nutrients in crop production. (in) Theme Papers on Integrated Nutrient Management. Kannaiyan S, Kumar K and Govindarajan K (Eds). Tamil Nadu Agriculture University and Tamil Nadu Department Agricultural, Coimbatore. 2000:87-101.
6. Duddigan S, Collins CD, Hussain Z, Osbahr H, Shaw LJ, Sinclair F, Sizmur T, Thallam V, Winowiecki LA. Impact of zero budget natural farming on crop yields in Andhra Pradesh, SE India. Sustainability. 2022;14(3):1689.
7. Palekar S. Text book on Shoonya Bandovalada naisargika Krushi, published by Swamy Anand, Agri Prakashana, Bangalore. 2006:170.
8. Anandan R, Priya L, Rajendran P. Dynamics of organic biofertilizers on *Oryza sativa* ADT43. International Journal of Current Microbiology and Applied Sciences. 2016;5(4):902-908.
9. Patel SK, Sharma A, Singh GS. Traditional agricultural practices in India: An approach for environmental sustainability and food security. Energy, Ecology and Environment. 2020;5(4):253–271.
10. Sharma IP, Kanta C, Dwivedi T, Rani R. Indigenous agricultural practices: A supreme key to maintaining biodiversity. In: Goel R, Soni R, Suyal D (eds) Microbiological advancements for higher altitude agro-ecosystems & sustainability Rhizosphere Biology. Springer Singapore; 2020.
11. Ravi V, Ravindran CS, Suja G, George J, Nedunchezhiyan M, Byju G, Naskar SK. Crop physiology of elephant foot yam [*Amorphophallus paeoniifolius* (Dennst. Nicolson)]. Adv. Hort. Sci. 2011;25(1):51-63.
12. Pramer D, Schmidt EL. Experimental soil microbiology. Minneapolis: Burgess Publ. Co; 1965.
13. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Indian Council of Agricultural Research Publication. New Delhi. 1985:87-89.
14. Bewley JD, Black M. Seeds: Physiology of development and germination. Springer Science & Business Media. 1985:332.
15. Lim J, Jee S, Lee D, Roh E, Jung K, Oh C, Heu S. Biocontrol of *Pectobacterium carotovorum* subsp. *carotovorum* using bacteriophage PP1. Journal of Microbiology and Biotechnology. 2013;23(8):1147–1153.
16. Chakraborty B, Sarkar I. Quality analysis and characterization of panchagavya, jeevumrutha and sasyamrutha. International Journal of Current Microbiology and Applied Sciences. 2019;8(5):2018-2026.
17. Gore NS, Sreenivasa MN. Influence of liquid organic manures on growth, nutrient content and yield of tomato (*Lycopersicon esculentum* Mill.) in the sterilized



- soil. Karnataka Journal of Agricultural Sciences. 2011;24(2):153-157.
18. Tao R, Liang YC, Wakelin SA, Chu GX. Supplementing chemical fertilizer with an organic component increases soil biological function and quality. *Applied Soil Ecology*. 2015;96:42–51.
  19. Lian TX, Wang GH, Yu ZH, Li YS, Liu XB, Zhang SQ, Herbert SJ, Jin J. Bacterial communities incorporating plant-derived carbon in the soybean rhizosphere in Mollisols that differ in soil organic carbon content. *Applied Soil Ecology*. 2017;119: 375-383.
  20. Biswas S, Jana K, Agrawal RK, Puste AM. Effect of integrated nutrient management on green forage, dry matter, and crude protein yield of oat in oat-lathyrus intercropping system. *Journal of Crop and Weed*. 2020;16(2):233-238.
  21. Mallick P. Amrut Jal for Plants: Preparation and Uses. In: *Organic Urban Farming, The Indian Way* (E-Book); 2016. Available:<http://organicterrace.in/blog/amrut-jal-organic-fertilizer-soil-conditioner/> Access on 30 December, 2022
  22. Kumar RP, Singh, ON, Yogeshwar S, Sachchidanand D, Singh JP. Effect of integrated nutrient management on growth, yield, nutrient uptake and economics of French bean (*Phaseolus vulgaris*). *Indian Journal of Agricultural Sciences*: 2009; 79(2):122-128.
  23. Mishra N. Growth and yield response of pea (*Pisum sativum* L.) to integrated nutrient management- a review. *Journal of Plant and Pest Science*. 2014;1(2):87-95.
  24. Kumar M, Chaudhary V, Naresh RK, Maurya OP, Pal SL. Does Integrated Sources of Nutrients Enhance Growth, Yield, Quality and Soil Fertility of Vegetable Crops? *International Journal of Current Microbiology and Applied Sciences*. 2018;7(6):125-155.
  25. Dutta AK, Mahato SK, Kumar K. Performance of organically grown french bean during post-kharif season in South Chhota Nagpur Plateau of Eastern India. *Journal of Bio Innovation*. 2022;11(1):33-43.
  26. Bhadu K, Sharma SK, Choudhary R, Yadav SK, Jain RK, Jat G, Ravishankar N. Effect of jeevamrut on productivity and profitability of wheat under organic farming. *Indian Journal of Agricultural Sciences*. 2021;91(6):924–928.
  27. Gopal V, Gurusiddappa LH. Influence of Jeevamrutha (fermented liquid manure) on growth and yield parameters of tomato (*Solanum Lycopersicum* L.). *World Journal of Environmental Biosciences*. 2022;11(3): 1-7.
  28. Yogananda SB, Thimmegowda P, Shruthi GK. Performance of cowpea [*Vigna unguiculata* (L.) Walp] under organic production system in southern dry zone of Karnataka. *Legume Research*. 2019;43(2): 229-234.
  29. Sreenivasa MN, Nagaraj N, Bhat SN. Beejamruth: A source for beneficial bacteria. *Karnataka Journal of Agricultural Sciences*. 2010;17(3):72-77.
  30. Dhanoji MM, Meena MK, Naik MC. Manure and plant extracts for foliar nutrition in organic farming: A Review. *International Journal of Chemical Studies*. 2018;6(6): 1447-1454.

© 2023 Biswas et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/97278>