



Effect of Different Doses of Irradiation on the Germination of Varieties of Maize Developing against the Fall Armyworm in the Central African Republic

Rufin Reo-Ndouba^a, Lucie Aba-Toumou^{a*},
Félix Allah-Barem^b, Augustin Doukofiona^b,
Steve Dieudonné Mbenda^b, Jephté Juste Kaïne^c,
Innocent Zinga^a, Silla Semballa^a
and Joseph Antoine Bell^d

^a Laboratory of Biological and Agronomical Sciences for Development, Faculty of Sciences, University of Bangui, Central African Republic.

^b Central African Institute of Agriculture Research, Central African Republic.

^c Higher Institute for Rural Development (ISDR), Mbaïki, University of Bangui, Central African Republic.

^d Genetics and Plant Breeding Unit, Faculty of Sciences, University of Yaoundé 1, Cameroon.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRCS/2023/v8i4179

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97387>

Original Research Article

Received: 03/02/2023

Accepted: 05/04/2023

Published: 17/04/2023

*Corresponding author: E-mail: rufincyrusrndb82@gmail.com;

ABSTRACT

Maize is a major crop feeding 80% of the population of the Central African Republic (CAF). However, the country is not yet self-sufficient in its maize need and continues to rely partially on imports. Exacerbating this situation is the recent invasive spread of the fall armyworm attributed to change in weather patterns since 2016 which has been decimating the crop leading to drastic yield losses. Various control measures continue to be tested globally for the control of FAW, including chemical control, biological control using microbial organisms and predatory insects that attack FAW, use of genetically modified crops with Bt genes, and integrated pest management. However, the use of developing genetic resistance in maize against the pest remains under-explored. We focused research on initiating a mutation breeding effort in maize in CAF.

The goal of this experience is to determinate the optimal dose from radio-sensitivity test of CMS8704 (100 Gy, 200 Gy, 300 Gy, 400 Gy and 500 Gy), CMS 2019 (100 Gy, 200 Gy, 300 Gy, 400 Gy and 500 Gy), CMS 8501 (100 Gy, 200 Gy, 300 Gy, 400 Gy and 500 Gy) and local ecotypes (100 Gy, 200 Gy, 300 Gy, 400 Gy and 500 Gy).

200 seeds are bulk to irradiation with different doses (100 Gy, 200 Gy, 300 Gy, 400 Gy and 500 Gy). The test of germination is carried out on 200 seeds four varieties. The treated seeds are sown at equal depths in a tray filled with soil/compost or can as well sow in natural condition on the field just to assure that the soil surface is flat, and the treated seeds are sown at equal depth containing the five treatments in rows of 200s seed each for one control and each treatment. Per assay three replicates are performed, one tray per replicate. Fourteen days after sowing, the seedling height and survival is measured to determine the Growth Reduction Value 50 or GR50.

All seeds from different doses (CMS-20 19, CMS87 04 and the Local ecotype) were germinated from different doses (100Gy, 200Gy and 300Gy) and did be presented the symptom of FAW. However, for the CMS85 01, all seeds from different doses (100Gy, 200Gy, 300Gy; 400Gy and 500Gy) were germinated as the control, but the major of plants are attacked by the FAW (Fall Armyworm).

Keywords: Host resistance; *Zea mays*; Fall Army Worm (FAW).

1. INTRODUCTION

“Maize (*Zea mays* L.) is a major cereal crop feeding 80% of the population of the Central African Republic (CAF)” [1]. “However, the country is not yet self-sufficient in its maize need and continues to rely partially on imports. Exacerbating this situation is the recent invasive spread of the Fall Armyworm (FAW) since 2016 which has been decimating the maize leading to drastic yield losses” [2]. The FAW attacks several crops of economic importance in CAF such as maize, rice, sorghum, millet, cowpea and peanut. The assessment of the FAW infestation situation in CAF showed an incidence ranging from 20% to 80% of attacked feet depending on the localities [2]. “The larval stages of the insect have about 300 plants hosts” [3]. “The ability of the adult stage to fly long distance of about 400 km per night, in part, accounts for their successful spread” [4]. “The most damaging FAW stage is larvae that feed on leaves, and lateral instars damage every part of the plant” [5]. “Due to its attack, the plant’s photosynthetic area

reduces, and it also directly damages the grains” [6-7].

Despite the importance of maize as an important food crop for CAF, its average productivity remains low in the country. Productivity is further reduced by the invasion of FAW.

“Various control measures continue to be tested globally for the control of FAW, including chemical control [8], biological control using microbial organisms and predatory insects that attack FAW [9], use of genetically modified crops with Bt genes, and integrated pest management” [10-12]. “However, the use of developing genetic resistance in maize against the pest remains under-explored” [13-15]. Induced genetic variation leads to a widened and novel genetic base for selections for traits of interest. This part of work focused on the identification of optimum dose of gamma irradiation treatment on seeds that could be used for improving productivity of maize and developing Resistance to the FAW using Radiation-Induced Novel Genetic Diversity.

2. MATERIALS AND METHODS

2.1. Choice Of Study Site

The experiment was carried out at the Regional Polyvalent Research Center (CRPR) of Boukoko (05°04' South latitude, 018°49' East longitude and 499 m altitude) in Region of Lobaye. It is a tropical type of climate marked by two main seasons (the rainy season from March to April and October to November and the dry season from J November/December to Feb [16].

The region has a strongly denatured, red ferrallitic soil [16]. The Boukoko region is covered by the subequatorial and peri-Guinean semi-deciduous forests. Following the strong anthropogenic pressure, some of these forests have disappeared and the initial vegetal landscape of the region has been modified. Thus, the forest is replaced by a vegetation of anthropic origin made up of spontaneous palm groves, grassy savannahs with *Panicum maximum*, *Imperata cylindrica*, and pre-forest savannahs often dominated by *Chromo-laena odorata* and *Caloconba Caloncaba weluitshii* [17].

2.2 Vegetable Materials

The test of performance from irradiated seeds was performed on three varieties of maize from Cameroon and on the local ecotype of CAR. The details of the characteristics of maize varieties used in this study are in Table 1.

2.3 Experimental Test And Measured Parameters

The experimental design adopted was the complete block plan with three repetitions (R1, R2 and R3) and six treatments (Controls, 100 Gy, 200 Gy, 300 Gy, 400 Gy and 500 Gy) [18]. There are four plots of 40 m long by 14 m wide for an area of 560 m² each. The repetitions were separated by 3 m in distance and 2 m of border at the ends. Sowing was carried out at spacings of 1 m on the line and 1.5 m between two lines at the rate of two seeds per pocket on May 07, 2021. A first application of inorganic manure (NPK) was carried out one day after the first weeding and the second application of inorganic manure (Urea) at the appearance of the male inflorescence. The measured parameters were in the four plots: (i) emergence rate (%), calculated two weeks after emergence; (ii) plant height (mm), measured with a tape measure at two

weeks and four weeks after sowing; (iii) plant survival (%) at two weeks and four weeks after sowing. The percentage of germination or emergence of seedlings (a), survival (b) and variation in seedling height (c), for each replication is calculated according to the following formula [19]:

- (a) Germination percentage = Number of germinated seeds / Number of seeds planted x 100
- (b) Survival percentage = Number of surviving seedlings / Number of seeds planted x 100
- (c) Average seedling height = Sum of seedling height in mm / Number of plants measured

2.4 Data Analysis

Statistical analysis of the data obtained was performed with Statistix software. The multicomparison Chi square test with the Bonferroni method was used to compare percentages of germination and survival at different doses (0Gy, 100 Gy; 200 Gy; 300 Gy; 400 Gy; 500 Gy). Due to inequalities in the numbers of survivals, the Generalized Linear Model (GLM) with error distribution of Poisson family was used to compare the averages of seedling heights. All tests were performed with the probability threshold of 5%. The graphics were generated using Excel software.

3. RESULTS

3.1 Data From Measured Parameters Cms 8704 Variety

Data from measured parameters CMS 8704 variety are reported in Table 2. with 100Gy and 200Gy all seeds were germinated as the control, with 300Gy and 400Gy the half of seed were germinated, with 500Gy 2 to 3 seed of each replicate were germinated, For all dose, there is not attack or symptom of FAW (Table 3).

3.2 Data from measured parameters CMS 2019 variety

All seeds from different doses (100Gy, 200Gy and 300Gy) were germinated (Table 2). With 400Gy the half of seed from each replicate were germinated, with 500Gy some replications give 2 to 3 plants. For all dose, there is not attack or symptom of FAW.

Table 1. Characteristics of maize varieties used in the study

Variety	Genetic nature (N Gq)	Origin & Year of release	Date of introduction or registration in CAR	Cycle (days)	Height of Plants (cm)	100 seed weight (g)	Colour of seed	Seed Texture	Potential Yield (t/ha)	Organoleptic characteristics and suitability for processing
CMS85 01	Composite	IRAD 1985)	1988	105-110	180-220	24.5	White	Cornea - dentate	5 - 8	Susceptibility to lodging, drought and stem borers
CMS-20 19	Composite	IRAD 1990)	1994	85-90	140-170	22, 8	White		4 - 5	Drought and disease tolerance, Striga ensitivity
CMS87 04	Composite	IRAD 1987)	1988	105-110	190-240	24.5	Yellow	Cornea	7 - 8	Sensitivity to lodging, good resistance to drought, very high sugar content

Local ecotype without information in the origine

Table 2. Data from measured parameters CMS 8704 variety

	Control	100 Gy	200 Gy	300 Gy	400 Gy	500 Gy
Germination (%)	100 ^a	93 ^a	92 ^a	60 ^b	28 ^c	2 ^d
Survival percentage (%)	100 ^a	90 ^a	95 ^a	38 ^b	17 ^c	0 ^d
N (survival)	60	54	57	23	10	0
Average seedling height (mean±SD)	203±14 ^a	194±11 ^a	199±13,5 ^a	168±12,3 ^b	105±10,7 ^c	-

Values with different letters are statistically different ($P<0,001$)

Table 3. Data from measured parameters CMS 2019 variety

	Control	100 Gy	200 Gy	300 Gy	400 Gy	500 Gy
Germination (%)	100 ^a	97 ^a	98 ^a	93 ^a	62 ^b	8 ^c
Survival percentage (%)	100 ^a	97 ^a	97 ^a	82 ^b	48 ^c	7 ^d
N (survival)	60	58	58	49	29	4
Average seedling height (mean±SD)	202±19 ^a	203±15 ^a	198±13.6 ^a	182±13,4 ^a	125±11 ^a	48±9,2 ^a

Values with different letters are statistically different ($P<0,001$)

Table 4. Data from measured parameters CMS 8501 variety

	Control	100 Gy	200 Gy	300 Gy	400 Gy	500 Gy
Germination (%)	100 ^a	93 ^a	95 ^a	95 ^a	65 ^b	30 ^c
Survival percentage (%)	95 ^a	90 ^a	85 ^a	87 ^a	55 ^b	22 ^c
N (survival)	57	54	51	52	33	13
Average seedling height (mean±SD)	202±15,2 ^a	202±14.5 ^a	201±12 ^a	191±12.3 ^a	137±7.8 ^b	81±6.5 ^c

Values with different letters are statistically different ($P<0,001$)

3.3 Data from Measured Parameters CMS 85 01 Variety

All seeds from diffrents doses (100Gy , 200Gy; 300Gy ; 400Gy and 500Gy) were germinated as the control, but the major of plants are attacked by the FAW (Table 4).

3.4 Data from Measured Parameters of Local Ecotype

With 100Gy and 200Gy all seeds were germinated as the control, but with hygh dose (300Gy ; 400Gy and 500Gy) the major of seeds were not germinated (Table 5) . We don't observe the symptom of FAW (Fall Armyworm).

4. DISCUSSION

"The use of ionizing irradiation has been useful for the genetic improvement crop. However, its use should be preceded by assays that determine the effect that irradiations have on the plant material. A radio stimulant low-dose is defined as any dose from environmental radiation level and the threshold that marks the boundary between positive and negative biological effect" [20]. "The effects of gamma radiation are investigated by studying plant germination, growth and development, and

biochemical characteristics of maize. Maize dry seeds are exposed to a gamma source at doses ranging from 0.1 to 1 kGy" [21-22]. Our results show that fir the CMS 8704 variety with 100Gy and 200Gy all seeds were germinated as the control, with 300Gy and 400Gy the half of seed were germinated, with 500Gy 2 to 3 seeds of each replicate were germinated, but according to the different doses, the CMS 8704 presents no symptom of FAW.

The data from measured parameters local ecotype show that with high dose (300Gy; 400Gy and 500Gy) the major of seeds were decreased plant germination and development. "These findings confirm the results obtained by earlier studies that showed the inhibitory effects of plant growth and development exposed to high doses of gamma radiation, simultaneous with the increase of reactive oxygen species generated through water radiolysis" [23-25]. "Massive doses of ionizing radiation have been shown to induce physiological changes in plants, such as enhancement of respiration, increase in ethylene production and induction of enzyme activities (particularly for phenolic metabolisms and accumulation specific protein species). These effects are considered a consequence of both the direct interactions between the ionizing radiation" [26-27].

Table 5. Data from measured parameters local ecotype

	Control	100 Gy	200 Gy	300 Gy	400 Gy	500 Gy
Germination (%)	100 ^a	180 ^a	140 ^b	50 ^c	25 ^d	0 ^d
Survival percentage (%)	195 ^a	155 ^a	130 ^b	45 ^c	15 ^d	-
N (survival)	40	31	26	9	3	0
Average seedling height (mean±SD)	202±10,8 ^a	202±12,2 ^a	201±10 ^a	164±7,4 ^b	83±8,1 ^c	-

Values with different letters are statistically different ($P < 0,001$).

5. CONCLUSION

All seeds from different doses (CMS-20 19, CMS8704 and the Local ecotype) were germinated from different doses (100Gy, 200Gy and 300Gy) and did be presented, the symptom of FAW. However for the CMS85 01, all seeds from different doses (100Gy, 200Gy; 300Gy; 400Gy and 500Gy) were germinated as the control, but the major of plants are attacked by the FAW (Fall Armyworm). Considering the effects of radiation on plants, the present study was conducted to determine the effects of radiation on maize growth and development and on the content of photosynthetic pigments. Furthermore, we used ESR spectroscopy to study the behavior of a radiation-induced free radical in gamma-irradiated maize seeds and to correlate it with the germination pattern.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Food and Agriculture Organization. National Strategy Document for the Integrated Management of the Fall Armyworm in the Central African Republic. 2019;41.
- Aba-Toumno L, et al. The first survey of *Spodoptera frugiperda* on Maize in the farms and in the traditional post-harvest conservation in Central African Republic. Asian Journal of Advances in Agriculture Research. 2018;8(3):1-6.
- Montezano DG, Specht A, Sosa-Gómez DR, et al. Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the America. Afr. Entomo; 2018. DOI: 10.4001/003.026.0286.
- Feldmann F, Rieckmann U, Winter S. The spread of the fall armyworm *Spodoptera frugiperda* in Africa—what should be done next? J. Plant Dis. Protect. 2019;126(2): 97-101
- Yu SJ, Nguyen SN, Abo-Elghar GE. Biochemical characteristics of insecticide resistance in the fall armyworm, *Spodoptera frugiperda* (J. E. Smith). Pesticide Biochemistry and Physiology 2003;77:1–11.
- Abrahams P, et al.. Fall Armyworm: Impacts and implications for Africa. Evidence note (2), September 2017. Report to DFID. Wallingford, UK: CAB International; 2017.
- Chimweta M, Nyakudya I, Jimu L, Mashigaidze AB. Fall armyworm *Spodoptera frugiperda* (JE Smith)] damage in maize: management options for flood-recession cropping smallholder farmers. Int. J. Pest Manage. 2020;66(2):142-154. Available:https://doi.org/10.1080/09670874.2019.1577514
- Chandrasena DI, Signorini AM, Abratti G, Storer NP, Olaciregui ML, Alves AP, Pilcher CD. Characterization of field-evolved resistance to *Bacillus thuringiensis*-derived Cry1F δ -endotoxin in *Spodoptera frugiperda* populations from Argentina. Pest Management Science Publié en line; 2017. DOI: 10.1002/ps.4776.
- Farias JR, Andow DA, Horikoshi RJ, Sorgatto RJ, Fresia P, Santos AC, Omoto C. Field-evolved resistance to Cry1F maize by *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Brazil. Crop Protection. 2014;64:150-158.
- Gerdes JT, Behr CF, Coors JG, Tracy WF. Compilation of North America maize breeding germplasm. Crop Science Society of America. 1993:202.
- Huang F, Qureshi JA, Meagher RL Jr, Reisig DD, Head GP, et al. Cry1F resistance in fall armyworm *Spodoptera frugiperda*: Single gene versus pyramided Bt maize. Plos One. 2014;9(11):e112958. DOI: 10.1371/journal.pone.0112958.
- Ni X, Xu W, Blanco MH, Williams PW. Evaluation of fall armyworm resistance in maize germplasm lines using visual leaf

- injury rating and predator survey. Insect Science. 2014;21:541-555.
13. Yu S, Nguyen S, Abo-Elghar G. Biochemical characteristics of insecticide resistance in the fall armyworm, *Spodoptera frugiperda* (JE Smith). Pesticide Biochemistry and Physiology. 2003;77(1):1-11.
14. Storer NP, Babcock JM, Schlenz M, Meade T, Thompson GD, Bing JW, Huckaba RM. Discovery and characterization of field resistance to Bt maize: *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Puerto Rico. Journal of Economic Entomology 2010;103:1031–1038.
15. Huang F, Qureshi JA, Meagher RL Jr, Reisig DD, Head GP, et al. Cry1F resistance in fall armyworm *Spodoptera frugiperda*: Single gene versus pyramided Bt maize. Plos One. 2014;9(11):e112958. DOI: 10.1371/journal.pone.0112958.
16. Boulvert Y. Carte phytogéographique de la République centrafricaine (feuille Ouest-feuille Est) à 1:1000000. ORSTOM. 1986;31.
17. Conway JL, Ouedraogo AK, Coneff J. Activité de zonage plus de moyens d'existence de la République centrafricaine. USAID (United States Agency International Development), Bangui, Centrafrique. 2012;41.
18. Spencer-Lopes, M.M. Forster, B.P. and Jankuloski, L. 2018. Manuel on mutation breeding. Plant Breeding and Genetics Subprogramme Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture Vienna, Austria, 3 ed, 319p.
19. Suresh D, Poonguzhali S, Sridharan S, Rajangam J. Determination of lethal dose for gamma rays induced mutagenesis in butter bean (*Phaseolus lunatus* L) Variety KKL-1. Int. J. Curr. Microbiol. App. Sci. 2017;6(3):712–717.
20. Iglesias-Andreu LG, Octavio-Aguilar P, Bello-Bello J. Current importance and potential use of low doses of gamma radiation in forest species. In Adrovic F (Ed.), Gamma radiation Rijeka, Croatia: In Tech Europe. Ikram N, Dawar S, Abbas Z. 2012:265–280.
21. Delia Marcu, Grigore Damian, Constantin Cosma, Victoria Cristea.. Gamma radiation effects on seed germination, growth and pigment content, and ESR study of induced free radicals in maize (*Zea mays*). J Biol Phys. 2013:625–634. DOI: 10.1007/s10867-013-9322-z
22. Weil CF, Monde RA. EMS mutagenesis and point mutation discovery. Molecular genetic approaches to maize improvement. Springer. 2009:161–171.
23. Strable J, Scanlon JM. Maize (*Zea mays*): a model organism for basic and applied research in plant biology. In: Emerging Model Organisms: A Laboratory Manual. Cold Spring Harbor Laboratory Press. 1999;2:33–41.
24. Esnault AM, Legue F, Chenal C. Ionizing radiation: Advances in plant response. Environ. Exp. Bot. 2010;68:231–237.
25. Dezfuli P, Sharif-Zadeh F, Janmohammadi M. Influence of priming techniques on seed germination behaviour of maize inbred lines (*Zea mays* L.). J. Agric. Biol. Sci. 2008;3:22–25.
26. Caplin N, Willey N. Ionizing radiation, higher plants, and radioprotection: From acute high doses to chronic low doses. Front. Plant Sci. 2018;9:847. [PubMed]
27. Zaka R, Chenal C, Misset MT. Study of external low irradiation dose effects on induction of chromosome aberrations in *Pisum sativum* root tip meristem. Mutat. Res. Genet. Toxicol. Environ. Mutagen. 2002;517:87–99.

© 2023 Reo-Ndouba et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/97387>